



Sutured Wound Care Instructions For Lips or Chin

- No strenuous activity for 48 hours. Resume moderate activity in 48 hours. No heavy exercising for one week.
- Take Extra Strength Tylenol (500 mg) or Acetaminophen as needed for discomfort, 1-2 tabs every 4-6 hours.
- Do not take any extra aspirin or anti-inflammatory products (Ibuprofen, Motrin, Advil, Aleve) for 1 week after surgery.
- Do not drink alcoholic beverages for 48 hours.
- Keep the white pressure bandage in place for 24 hours. If the bandage becomes blood tinged or loose, reinforce it with gauze and tape. (See reverse side for management of bleeding). Remove the white bandage in 24 hours
- Leave the flat tan bandage in place for 1 week. Keep the bandage dry. Wash around it carefully.
- If the tape becomes soiled or starts to come off, reinforce it with additional paper tape.
- Do not smoke for 3 weeks; smoking is detrimental to wound healing.
- It is normal to have swelling and bruising around the surgical site. The bruising will fade in approximately 10-14 days. Elevate the area to reduce swelling.
- Try to keep your lips / chin as immobile as possible. Refrain from laughing, smiling and yawning for 3 weeks. Eat soft foods for the first 24 hours and take small bites of food for the entire three weeks.
- When brushing your teeth, you should use a child's toothbrush or use mouth wash to prevent stretching of surgery site.
- Numbness, itchiness and sensitivity to temperature changes can occur after surgery and may take up to 18 months to normalize.

➤ **IN CASE OF EMERGENCY: 317-859-8970**

INSTRUCTIONS FOR SECOND WEEK:

- Remove the tan bandage and steri-strips
- Clean incision **gently** with tap water and mild soap, using a Q-tip, cotton ball or gauze (shower/bathe normally). You may use a small amount of Peroxide to gently remove any scabbed areas.
- Apply Vaseline or Aquaphor with a Q-tip (Do **NOT** use Neosporin or Triple Antibiotic ointment). It is OK to use Bacitracin or Polysporin for 1-2 weeks.
- Cover the incision with a bandaid or nonstick gauze and paper tape.
- Repeat **daily** for one week or until completely healed with no open or draining areas.

POSSIBLE COMPLICATIONS

BLEEDING:

1. Leave the bandage in place.
2. Use tightly rolled up gauze or a cloth to apply direct pressure over the bandage for 20 minutes.
3. Reapply pressure for an additional 20 minutes if necessary
4. Call the office or go to the nearest emergency room if pressure fails to stop the bleeding.
5. Use additional gauze and tape to maintain pressure once the bleeding has stopped.

PAIN:

1. Post operative pain should slowly get better, never worse.
2. A severe increase in pain may indicate a problem. Call the office if this occurs.

INFECTION:

1. Watch for fevers, discharge, spreading redness, increased tenderness, usually 4-7 days after the procedure. Call the office if this occurs.

GENERAL INFORMATION REGARDING YOUR INCISION

*Once the bandages are removed, the scar will be red and firm (especially in the lip/chin area). This is normal and will fade in time. It might take 6-12 months for this to happen.

*Massaging the area will help the scar soften and fade quicker. Begin massage **one month** after the bandages have been removed. Apply pressure directly and firmly over the scar with the fingertips and move in a circular motion. Massage the area for a few minutes several times a day.

*6-8 weeks after surgery it is not uncommon to see the formation of “tender pimple-like” bump along the scar. This is normal. As the scar continues to mature and the stitches underneath the skin begin to dissolve, this might occur. Do not pick or squeeze, this will resolve on its own. Should one break open producing a small amount of drainage, apply Polysporin or Bacitracin ointment a few times a day until the wound is completely healed.

*Numbness in the surgical area is expected. It might take 12-18 months for the feeling to return to normal. During this time sensations of itchiness, tingling and occasional sharp pains might be noted. These feelings are normal and will subside once the nerves have completely healed.

***Protect a fresh scar from the sun.** Use a sun-screen with SPF 30 or greater or barriers as clothing, or a wide-brimmed hat.